

BMP-4/BMP-2B/DVR4, Human, Recombinant

货号：PCK269

产品信息

别名	BMP-2B, DVR4
物种	Human
表达宿主	E.coli
序列信息	MKKNKNCRRHSLYVDFSDVGWNDWIVAPPGYQAFYCHGDCPFPLADHLN STNHAIVQTLVNSVNSSIPKACCVPTELSAISMLYLDEYDKVVLKKNYQEMV VEGCGCR with polyhistidine tag at the C-terminus.
检索号	P12644.1
分子量	12.88 kDa
标签	His-tag at the C-terminus
生物活性	Measure by its ability to induce alkaline phosphatase production by ATDC5 cells. The ED50 for this effect is <0.58 ng/mL.

产品特性

纯度	>95% as determined by SDS-PAGE. Ni-NTA chromatography.
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 20 mM sodium carbonate, pH 9.0.



复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 $\mu\text{g/mL}$. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Bone morphogenetic protein 4 is a protein that in humans is encoded by BMP4 gene. BMP4 is found on chromosome 14q22-q23. BMP4 is a member of the bone morphogenetic protein family which is part of the transforming growth factor-beta superfamily. The superfamily includes large families of growth and differentiation factors. BMP4 is highly conserved evolutionarily. BMP4 is found in early embryonic development in the ventral marginal zone and in the eye, heart blood and otic vesicle.

SDS-PAGE

