

## Pleiotrophin/PTN/HARP/HBGF-8/OSF-1, Human, Recombinant

货号：PCK297

### 产品信息

别名	PTN, Heparin Affin Regulatory Protein (HARP), Heparin-Binding Growth Factor-8 (HBGF-8), Osteoblast-Specific Factor 1 (OSF-1)
物种	Human
表达宿主	E.coli
序列信息	MGKKEKPEKKVKKSDCGEWQWSVCVPTSGDCGLGTREGTRTGAECKQT MKTQRCKIPCNWKKQFGAECKYQFQAWGECDLNTALKTRTGSLKRALHN AECQKTVTISKPCGKLTKPKPQAESKKKKKEGKKQEKMLD with polyhistidine tag at the C-terminus.
检索号	P21246.1
分子量	19.89 kDa
标签	His-tag at the C-terminus

### 产品特性

纯度	>98% as determined by SDS-PAGE. Ni-NTA chromatography
内毒素	<0.01 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.



## 复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

## 背景介绍

Pleiotrophin (PTN) also known HBBM, HBGF-8, NEGF1, HARP or HB-GAM is a protein that in humans is encoded by the PTN gene. This protein is expressed in the central and peripheral nervous system and also in several non-neural tissues, notably lung, kidney, gut and bone. Pleiotrophin binds anaplastic lymphoma kinase (ALK) which induces MAPK pathway activation, an important step in the anti-apoptotic signaling of PTN and regulation of cell proliferation.

## SDS-PAGE

