

## M-CSF/CSF-1/MGI-IM, Mouse, Recombinant

货号 : PCK303

### 产品信息

别名	CSF-1, MGI-IM
物种	Mouse
表达宿主	E.coli
序列信息	MKEVSEHC SHMIGNGHLKVLQQLIDSQMETSCQIAFEFVDQEQLDDPVCYL KKAFFLVQDIIDETMRFKDNTPNANATERLQELSNNLNSCFTKDYEEQNKA CVRTFHETPLQLLEKIKNFFNETKNLLEKDWNIFTKNCNNSFAKCSSRDVVT KP with polyhistidine tag at the C-terminus
检索号	Q9QYH9.1
分子量	19.02 kDa
标签	His-tag at the C-terminus
生物活性	Measure by its ability to induce proliferation in NFS-60 cells. The ED50 for this effect is <2 ng/mL. The specific activity of recombinant mouse M-CSF is approximately >5 x 10 <sup>5</sup> IU/mg.

### 产品特性

纯度	>98% as determined by SDS-PAGE. Ni-NTA chromatography
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0.



## 复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

## 背景介绍

Macrophage Colony-Stimulating Factor (M-CSF), is a secreted cytokine which causes hematopoietic stem cells to differentiate into macrophages or other related cell types. The active form of M-CSF/CSF-1 is found extracellularly as a disulfide-linked homodimer, and is thought to be produced by proteolytic cleavage of membrane-bound precursors. M-CSF/CSF-1 induces cells of the monocyte/macrophage lineage. It also plays a role in immunological defenses, bone metabolism, lipoproteins clearance, fertility and pregnancy. Upregulation of M-CSF/CSF-1 in the infarcted myocardium may have an active role in healing not only through its effects on cells of monocyte/macrophage lineage, but also by regulating endothelial cell chemokine expression.

## SDS-PAGE

