

IL-18/IGIF, Mouse, Recombinant

货号: PCK311

产品信息

别名 **IGIF**

物种 Mouse

表达宿主 E.coli

序列信息 MNFGRLHCTTAVIRNINDQVLFVDKRQPVFEDMTDIDQSASEPQTRLIIYMY

> KDSEVRGLAVTLSVKDSKMSTLSCKNKIISFEEMDPPENIDDIOSDLIFFOKR VPGHNKMEFESSLYEGHFLACQKEDDAFKLILKKKDENGDKSVMFTLTNL

HQS with polyhistidine tag at the C-terminus.

检索号 P70380.2

分子量 19.1 kDa

标签 His-tag at the C-terminus

生物活性 Measure by its ability to induce IFN gamma secretion in KG-1 cells. The ED50 for

this effect is $<0.5 \mu g/mL$.

产品特性

>98% as determined by SDS-PAGE. Ni-NTA chromatography 纯度

内毒素 <0.1 EU per 1 µg of the protein by the LAL method.

保存 Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution,

> store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g.0.1%BSA,10%FBS,5%HSA or 5% trehalose solution),

protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.

运输 Ambient temperature or ice pack.

制剂 The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH

8.0.

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋








复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μ g/mL.Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Interleukin-18 (L-18) is a cytokine that belongs to the IL-1 superfamily and is produced by macrophages and other cells. IL-18 works by binding to the interleukin-18 receptor, and together with IL-12 it induces cell-mediated immunity following infection with microbial products like lipopolysaccharide (LPS). After stimulation with IL-18, natural killer (NK) cells and certain T cells release another important cytokine called interferon-γ (IFN-γ) or type II interferon that plays an important role in activating the macrophages or other cells. The combination of this cytokine and IL12 has been shown to inhibit IL-4 dependent IgE and IgG1 production, and enhance IgG2a production in B cells. IL-18 binding protein (IL18BP) can specifically interact with this cytokine, and thus negatively regulate its biological activity.

SDS-PAGE



普诺赛® | Pricella

网站: <u>www.procell.com.cn</u> 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



