

## IL-3/HCGFBPA/PSF, Mouse, Recombinant

货号 : PCK312

### 产品信息

别名	MCGF (Mast Cell Growth Factor), Multi-CSF, HCGF, P-cell stimulation factor, Interleukin-3b, BPA, PSF, Csfmu
物种	Mouse
表达宿主	E.coli
序列信息	MDTHRLTRTLNCSSIVKEIIGKLPPELKTDDDEGPSLRNKSFRRVNLSKRFVES QGEVDPEDRYVIKSNLQKLNCCCLPTSANDSALPGVFIRDLDLDDFRKKLRFYM VHLNDLETVLTSRPPQPASGSVSPNRGTVEC with polyhistidine tag at the C-terminus.
检索号	P01586.1
分子量	16.04 kDa
标签	His-tag at the C-terminus
生物活性	Measure by its ability to induce NFS-60 cells proliferation. The ED50 for this effect is <85 pg/mL. The specific activity of recombinant mouse IL-3 is approximately >1x 10 <sup>7</sup> IU/mg.

### 产品特性

纯度	>98% as determined by SDS-PAGE. Ni-NTA chromatography
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.



## 复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

## 背景介绍

Interleukin 3 is an interleukin, a type of biological signal (cytokine) that can improve the body's natural response to disease as part of the immune system. It acts by binding to the interleukin-3 receptor. Interleukin 3 stimulates the differentiation of multipotent hematopoietic stem cells into myeloid progenitor cells or, with the addition of IL-7, into lymphoid progenitor cells. In addition, IL-3 stimulates proliferation of all cells in the myeloid lineage (granulocytes, monocytes, and dendritic cells), in conjunction with other cytokines, e.g., Erythropoietin (EPO), Granulocyte macrophage colony-stimulating factor (GM-CSF), and IL-6. It is secreted by basophils and activated T cells to support growth and differentiation of T cells from the bone marrow in an immune response.

## SDS-PAGE

