

IL-4/IL4, Swine, Recombinant

货号：PCK343

产品信息

别名	IL4
物种	Swine
表达宿主	E.coli
序列信息	MHKCDITLQEIIKTLNILTARKNSCMELPVTDVFAAPENTTEKETFCRASTV LRHIYRHHTCMKSLLSGLDRNLSSMANMTCSVHEAKKSTLKDFLERLKTIM KEKYSKC with polyhistidine tag at the C-terminus.
检索号	Q04745.1
分子量	13.5 kDa
标签	His-tag at the C-terminus
生物活性	Measure by its ability to induce proliferation in TF-1 cells. The ED50 for this effect is <0.6 ng/mL.

产品特性

纯度	>98% as determined by SDS-PAGE. Ni-NTA chromatography
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.



复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 $\mu\text{g/mL}$. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

The interleukin 4 (IL4, IL-4) is a cytokine that induces differentiation of naive helper T cells (Th0 cells) to Th2 cells. Upon activation by IL-4, Th2 cells subsequently produce additional IL-4 in a positive feedback loop. The cell that initially produces IL-4, thus inducing Th2 differentiation, has not been identified, but recent studies suggest that basophils may be the effector cell. It is closely related and has functions similar to Interleukin 13.

SDS-PAGE

