

EGF/Urogastrone/URG, Swine, Recombinant

货号: PCK348

产品信息

别名 Urogastrone, URG

物种 Swine

表达宿主 E.coli

ACQHRE Pricella 序列信息 MNSYSECPPSHDGYCLHGGVCMYIEAVDSYACNCVFGYVGERCQHRDLK

WWELR with polyhistidine tag at the C-terminus.

检索号 Q00968.2

分子量 7.1 kDa

标签 His-tag at the C-terminus

产品特性

纯度 >95% as determined by SDS-PAGE. Ni-NTA chromatography.

内毒素 <0.1 EU per 1 µg of the protein by the LAL method.

保存 Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution,

> store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g.0.1%BSA,10%FBS,5%HSA or 5% trehalose solution),

protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.

运输 Ambient temperature or ice pack.

制剂 The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH

8.0.

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋








复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL.Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Epidermal growth factor (EGF) stimulates cell growth and differentiation by binding to its receptor, EGFR. Human EGF is a 6-kDa protein with 53 amino acid residues and three intramolecular disulfide bonds. EGF is present in various body fluids, including blood, milk, urine, saliva, seminal fluid, pancreatic juice, cerebrospinal fluid, and amniotic fluid. Biological activities ascribed to EGF include epithelial development, angiogenesis, inhibition of gastric acid secretion, fibroblast proliferation, and colony formation of epidermal cells in culture.

SDS-PAGE



网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



