

IL-2 (C145S), Human, Recombinant

货号 : PCK021

产品信息

别名	Interleukin-2; IL-2; T-cell Growth Factor; TCGF; Aldesleukin
物种	Human
表达宿主	E.coli
序列信息	Pro22-Thr153(Cys145Ser)
检索号	P60568
分子量	15.5 kDa
生物活性	Measured in a cell proliferation assay using CTLL-2 mouse cytotoxic T cells. The ED50 for this effect is 90-270 pg/ml.

产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	<1.0 EU per µg as determined by LAL test.
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 µm filtered solution of 10mM Acetata-Na, 5% Trehaiose, pH 4.5.




复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


背景介绍

Recombinant Human Interleukin-2 is a highly purified protein with a molecular weight of approximately 15,300 Daltons. The chemical name is des-alanyl-1, serine-145 Human Interleukin-2. It is produced by recombinant DNA technology using a genetically engineered E. coli strain containing an analog of the human interleukin-2 gene. Genetic engineering techniques were used to modify the Human IL-2 gene, and the resulting expression clone encodes a modified Human IL-2. This recombinant form differs from native Interleukin-2 in following ways: it is not glycosylated; the molecule has serine substituted for cysteine at amino acid position 145; the aggregation state of molecule is likely to be different from that of native IL-2.

SDS-PAGE



生物活性



Measured in a cell proliferation assay using CTLL-2 mouse cytotoxic T cells. The ED50 for this effect is 90-270 pg/ml.

