

IL-4/ BSF-1 (C-6His), Mouse, Recombinant

货号: PCK034

产品信息

别名	Interleukin-4; IL-4; IL4; B-cell IgG differentiation factor; B-cell Growth Factor 1;
	B-cell stimulatory factor 1; BSF-1; IGG1 induction factor; Lymphocyte stimulatory
	factor 1
物种	Mouse
表达宿主	Human Cells
序列信息	His21-Ser140
检索号	Human Cells His21-Ser140 P07750
分子量	14.6 kDa
标签	C-6His
生物活性	Measured in a cell proliferation assay using M-NFS-60 mouse lymphoblast cells. The ED50 for this effect is 43.59pg/ml.
产品特性	普诺廷 Procell

产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	<1.0 EU per µg as determined by LAL test.
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.





复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml.Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

Interleukin-4 (IL-4) is a pleiotropic Cytokine that regulates diverse T and B cell responses including cell proliferation, survival and gene expression. IL-4 is produced by mast cells, T cells, and bone marrow stromal cells. IL-4 regulates the differentiation of naive CD4+ T cells into helper Th2 cells, characterized by their Cytokine-secretion profile that includes secretion of IL-4, IL-5, IL-6, IL-10, and IL-13, which favor a humoral immune response. Another dominant function of IL-4 is the regulation of immunoglobulin class switching to the IgG1 and IgE isotypes. Excessive IL-4 production by Th2 cells has been associated with elevated IgE production and allergic response.



