

IL-4/ BSF-1 (C-6His), Mouse, Recombinant

货号：PCK034

产品信息

| | |
|------|---|
| 别名 | Interleukin-4; IL-4; IL4; B-cell IgG differentiation factor; B-cell Growth Factor 1; B-cell stimulatory factor 1; BSF-1; IGG1 induction factor; Lymphocyte stimulatory factor 1 |
| 物种 | Mouse |
| 表达宿主 | Human Cells |
| 序列信息 | His21-Ser140 |
| 检索号 | P07750 |
| 分子量 | 14.6 kDa |
| 标签 | C-6His |
| 生物活性 | Measured in a cell proliferation assay using M-NFS-60 mouse lymphoblast cells. The ED50 for this effect is 43.59pg/ml. |

产品特性

| | |
|-----|---|
| 纯度 | >95% as determined by reducing SDS-PAGE. |
| 内毒素 | <1.0 EU per µg as determined by LAL test. |
| 保存 | Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months. |
| 运输 | Ambient temperature or ice pack. |
| 制剂 | Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. |




复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


背景介绍

Interleukin-4 (IL-4) is a pleiotropic Cytokine that regulates diverse T and B cell responses including cell proliferation, survival and gene expression. IL-4 is produced by mast cells, T cells, and bone marrow stromal cells. IL-4 regulates the differentiation of naive CD4+ T cells into helper Th2 cells, characterized by their Cytokine-secretion profile that includes secretion of IL-4, IL-5, IL-6, IL-10, and IL-13, which favor a humoral immune response. Another dominant function of IL-4 is the regulation of immunoglobulin class switching to the IgG1 and IgE isotypes. Excessive IL-4 production by Th2 cells has been associated with elevated IgE production and allergic response.

SDS-PAGE



生物活性



Measured in a cell proliferation assay using M-NFS-60 mouse lymphoblast cells. The ED50 for this effect is 43.59pg/ml.

