

IL-1β/IL1F2, Human, Recombinant

货号: PCK047

产品信息

别名 Catabolin, Lymphocyte-Activating Factor (LAF), Endogenous Pyrogen (EP),

Leukocyte Endoge_x0002_nous Mediator (LEM), Mononuclear Cell Factor (MCF)

物种 Human

表达宿主 E.coli

序列信息 MASAPVRSLNCTLRDSQQKSLVMSGPYELKALHLQGQDMEQQVVFSMSFV

> QGEESNDKIPVALGLKEKNLYLSCVLKDDKPTLQLESVDPKNYPKKKMEK RFVFNKIEINNKLEFESAQFPNWYISTSQAENMPVFLGGTKGGQDITDFTMQ

FVSS with polyhistidine tag at the C-terminus.

检索号 P01584.2

分子量 18.48 kDa

标签 His-tag at the C-terminus

生物活性 Measure by its ability to induce proliferation in D10.G4.1 cells. The ED50 for this

> effect is <10 pg/mL. The specific activity of recombinant human IL-1 beta is approximately >1.5 x 10⁸ IU/mg. Measure by its ability to induce IL-8 secretion in -cretic

HT29 cells. The ED50 for this effect is 1.8-5.1 ng/mL.

产品特性

纯度 >98% as determined by SDS-PAGE. Ni-NTA chromatography

内毒素 <0.1 EU per 1 µg of the protein by the LAL method.

保存 Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution,

> store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g.0.1% BSA,10%FBS,5%HSA or 5% trehalose solution),

protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.

运输 Ambient temperature or ice pack.

制剂 The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH

8.0.

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋







复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL.Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Interleukin 1 beta (IL-1β) also known as leukocytic pyrogen, leukocytic endogenous mediator, mononuclear cell factor, lymphocyte activating factor and other names, is a cytokine protein that in humans is encoded by the IL1B gene. There are two genes for interleukin-1 (IL-1): IL-1 alpha and 遊講蹇® | Pricella IL-1 beta (this gene). IL-1β precursor is cleaved by cytosolic caspase 1 (interleukin 1 beta convertase) to form mature IL-1β.

SDS-PAGE

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



