

TNF- α / TNFA/ TNFSF2 (N-6His), Human, Recombinant

货号：PCK052

产品信息

别名	Tumor Necrosis Factor; Cachectin; TNF-Alpha; Tumor Necrosis Factor Ligand Superfamily Member 2; TNF-a; TNF; TNFA; TNFSF2
物种	Human
表达宿主	E.coli
序列信息	Gly57-Leu233
检索号	P01375
分子量	21.8 kDa
标签	N-6His
生物活性	Measured in a cytotoxicity assay using L-929 mouse fibroblast cells in the presence of the metabolic inhibitor actinomycin D. The ED50 for this effect is 30-150 pg/ml.

产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	<1.0 EU per μ g as determined by LAL test.
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 μ m filtered solution of 20mM PB, 100mM NaCl, pH 7.2.




复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


背景介绍

Tumor Necrosis Factor- α (TNF- α) is secreted by macrophages, monocytes, neutrophils, T-cells, and NK-cells following stimulation by bacterial LPS. Cells expressing CD4 secrete TNF- α while cells that express CD8 secrete little or no TNF- α . Synthesis of TNF- α can be induced by many different stimuli including interferons, IL2, and GM-CSF. The clinical use of the potent anti-tumor activity of TNF- α has been limited by the proinflammatory side effects such as fever, dose-limiting hypotension, hepatotoxicity, intravascular thrombosis, and hemorrhage. Designing clinically applicable TNF- α mutants with low systemic toxicity has been of intense pharmacological interest. Human TNF- α that binds to murine TNF-R55 but not murine TNF-R7, exhibits retained anti-tumor activity and reduced systemic toxicity in mice compared with murine TNF- α , which binds to both murine TNF Receptors. Based on these results, many TNF- α mutants that selectively bind to TNF-R55 have been designed. These mutants displayed cytotoxic activities on tumor cell lines in vitro and have exhibited lower systemic toxicity in vivo. Recombinant Human TNF- α High Active Mutant differs from the wild-type by amino acid substitution of amino acids 1-7 with Arg8, Lys9, Arg10 and Phe157. This mutant form has been shown to have increased activity with less inflammatory side effects in vivo.

SDS-PAGE



生物活性



Measured in a cytotoxicity assay using L-929 mouse fibroblast cells in the presence of the metabolic inhibitor actinomycin D. The ED50 for this effect is 30-150 pg/ml.

