

CCL2/MCP-1, Human, Recombinant

货号 : PCK111

产品信息

别名	Monocyte Chemotactic Protein-1, MCP-1
物种	Human
表达宿主	E.coli
序列信息	QPDAINAPVTCCYNFTNRKISVQRLASYRRITSSKCPKEAVIFKTIVAKEICA DPKQKWVQDSMDHLDKQTQTPKT with polyhistidine tag at the N-terminus.
检索号	P13500.1
分子量	9.49 kDa
标签	His-tag at the N-terminus
生物活性	Measure by its ability to chemoattract BaF3 cells transfected with human CCR2A. The ED50 for this effect is <20 ng/mL.

产品特性

纯度	>95% as determined by SDS-PAGE. Ni-NTA chromatography.
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.



复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

CCL2, also known as MCP-1, is belonging to the CC β chemokine family. CCL2 can be identified in endothelial cells, smooth muscle cells and monocytes as the results of reaction to several atherogenic stimulants, such as CD40 ligand, (IL-1β) and oxidized low density lipoprotein, interleukin-1β platelet derived growth factor (PDGF). Recent study shows that in vivo MCP1 have several critical roles in atherosclerosis. Additionally, MCP-1 has been proved involving in monocytic infiltration of tissues during several inflammatory diseases, and has been implicated in macrophage-mediated tumor growth inhibition in mice. In addition, CCL2 has been shown to have direct effects on tumor cells in an autocrine and paracrine fashion in multiple cancers, including sarcoma, lung, cervix, ovary, breast, and prostate.

SDS-PAGE

