

RANTES/ CCL5, Human, Recombinant

货号 : PCK114

产品信息

别名	C-C Motif Chemokine 5; EoCP; Eosinophil Chemotactic Cytokine; SIS-Delta; Small-Inducible Cytokine A5; T Cell-Specific Protein P228; TCP228; T-Cell-Specific Protein RANTES; CCL5; D17S136E; SCYA5
物种	Human
表达宿主	E.coli
序列信息	Ser24-Ser91
检索号	P13501
分子量	7.8 kDa

产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	<1.0 EU per µg as determined by LAL test.
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 µm filtered solution of 20mM Citrate, 6% Trehalose, 4% Mannitol, 0.05% Tween 80, pH4.0.




复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

Human Chemokine (C-C Motif) Ligand 5 (CCL5) plays an active role in recruiting leukocytes into inflammatory sites. CCL5 is secreted by many cell types at inflammatory sites and it exerts a wide range of activities through the Receptors CCR1, CCR3, CCR4, and CCR5. N-Terminal truncated CCL5/ RANTES, Met-RANTES, and amino-oxypentane (AOP)-RANTES exhibit antagonist or partial agonist functions on their Receptors. CCL5/ RANTES attracts different subtypes of leukocytes into inflamed tissue and intervenes in a wide range of allergic and autoimmune diseases.

SDS-PAGE



www.procell.com.cn

