

GRO-y/CXCL3 (N-6His), Human, Recombinant

货号: PCK125

产品信息

别名 C-X-C Motif Chemokine 3; GRO-Gamma (1-73); Growth-Regulated Protein

Gamma; GRO-Gamma; Macrophage Inflammatory Protein 2-Beta; MIP2-Beta;

GRO-Gamma (5-73); CXCL3; GRO3; GROG; SCYB3

物种 Human

表达宿主 E.coli

序列信息 Ala35-Asn107

检索号 P19876

分子量 10.1 kDa

标签 N-6His

产品特性

纯度 >95% as determined by reducing SDS-PAGE.

内毒素 <1.0 EU per µg as determined by LAL test.

保存 Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt.

Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at -5~-20°C for 3 months.

运输 Ambient temperature or ice pack.

制剂 Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋








复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml.Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

C-X-C Motif Chemokine 3 (CXCL3) is a secreted Protein that belongs to the intercrine alpha (Chemokine CXC) family. CXCL3 controls the migration and adhesion of monocytes and mediates its effect on its target cell by interacting with a cell surface Chemokine Receptor called CXCR2. In addition, CXCL3 is thought to play a role in inflammation and exert its effects on endothelial cells in an autocrine fashion.

SDS-PAGE



网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



