

IL-18RAcP/ IL-1R7 (C-6His), Human, Recombinant

货号: PCK150

产品信息

别名 Interleukin-18 Receptor accessory Protein (IL18RAP); Accessory Protein-like;

CD218 antigen-like family member B; CDw218b; IL-1R accessory Protein-like;

Interleukin-1 Receptor 7; Interleukin-18 Receptor accessory Protein-like;

Interleukin-18 Receptor

物种 Human

表达宿主 **Human Cells**

序列信息 Phe20-Arg356

检索号 O95256

分子量 39.4 kDa

标签 C-6His

产品特性

纯度 >95% as determined by reducing SDS-PAGE.

内毒素 <1.0 EU per µg as determined by LAL test.

保存 Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt.

Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at -5~-20°C for 3 months.

运输 Ambient temperature or ice pack.

制剂 Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋








复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml.Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

IL-18RAP is a single-pass type I membrane Protein and contains two Ig-like C2-type domains and one TIR domain, IL18RAP can be induced by IFN-alpha and IL12 in nature killer cells and T-cells. and JN pricella The coexpression of IL18R1 and IL18RAP is required for the activation of NF-kappa B and JNK in response to IL-18.Defects in IL18RAP are associated with Coeliac disease.

SDS-PAGE



网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



