

IL-18/IGIF/IL-1g/IL1F4, Human, Recombinant

货号 : PCK179

产品信息

别名 IGIF, IL-1g, IL1F4

物种 Human

表达宿主 E.coli

序列信息 MYFGKLESKLSVIRNLNDQVLFIDQGNRPLFEDMTDSDCRDNAPRTIFIISM
YKDSQPRGMAVTISVKCEKISTLSCENKIISFKEMNPPDNIKDTKSDIIFQRS
VPGHDNKMQFESSYEGYFLACEKERDLFKLILKKEDELGDRSIMFTVQNE
D with polyhistidine tag at the C-terminus.

检索号 Q14116.1

分子量 18.2 kDa

标签 His-tag at the C-terminus

生物活性 Measure by its ability to induce IFN gamma secretion in KG-1 cells. The ED50 for this effect is <5 ng/mL.

产品特性

纯度 >98% as determined by SDS-PAGE. Ni-NTA chromatography

内毒素 <0.1 EU per 1 µg of the protein by the LAL method.

保存 Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g.0.1% BSA,10%FBS,5%HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.

运输 Ambient temperature or ice pack.

制剂 The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 8.0.



复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 $\mu\text{g/mL}$. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Interleukin-18 (IL-18) is a cytokine that belongs to the IL-1 superfamily and is produced by macrophages and other cells. IL-18 works by binding to the interleukin-18 receptor, and together with IL-12 it induces cell-mediated immunity following infection with microbial products like lipopolysaccharide (LPS). After stimulation with IL-18, natural killer (NK) cells and certain T cells release another important cytokine called interferon- γ (IFN- γ) or type II interferon that plays an important role in activating the macrophages or other cells. The combination of this cytokine and IL12 has been shown to inhibit IL-4 dependent IgE and IgG1 production, and enhance IgG2a production in B cells. IL-18 binding protein (IL18BP) can specifically interact with this cytokine, and thus negatively regulate its biological activity.

SDS-PAGE

