

IL-22 (C-hIgG4 Fc), Human, Recombinant

货号：PCK186

产品信息

别名	Interleukin-22; IL-22; Cytokine Zcyto18; IL-10-related T-cell-Derived-inducible factor; IL-TIF; IL22
物种	Human
表达宿主	Human Cells
序列信息	Ala34-Ile179
检索号	Q9GZX6
分子量	43.4 kDa
标签	C-hIgG4 Fc

产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	<1.0 EU per μg as determined by LAL test.
保存	Lyophilized protein should be stored at $-5\sim-20^{\circ}\text{C}$, stable for one year after receipt. Reconstituted protein solution can be stored at $2-8^{\circ}\text{C}$ for 2-7 days. Aliquots of reconstituted samples are stable at $-5\sim-20^{\circ}\text{C}$ for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a $0.2\ \mu\text{m}$ filtered solution of PBS, pH7.4.



复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

Interleukin-22 (IL-22) is a member of a group of the IL-10 family, a class of potent mediators of cellular inflammatory responses. IL-22 is produced by activated DC and T cells. IL-22 and IL-10 Receptor chains play a role in cellular targeting and signal transduction. It can initiate and regulate innate immune responses against bacterial pathogens especially in epithelial cells such as respiratory and gut epithelial cells. IL-22 along with IL-17 likely plays a role in the coordinated response of both adaptive and innate immune systems. IL-22 also promotes hepatocyte survival in the liver and epithelial cells in the lung and gut similar to IL-10. Biological activity of IL-22 is initiated by binding to a cell-surface complex consisting of IL-22R1 and IL-10R2 Receptor chains. IL-22 biological activity is further regulated by interactions with a soluble binding Protein, IL-22BP. IL-22BP and an extracellular region of IL-22R1 share sequence similarity. In some cases, the pro-inflammatory versus tissue-protective functions of IL-22 are regulated by Cytokine IL-17A.

SDS-PAGE

