

## MCP-1/ CCL2 (C-6His), Mouse, Recombinant

货号：PCK228

### 产品信息

别名	C-C Motif Chemokine 2; Monocyte chemoattractant Protein 1; Monocyte chemotactic Protein 1; MCP-1; Platelet-Derived Growth Factor-inducible Protein 1; Small-inducible Cytokine A2; Ccl2; Je; Mcp1; Scya2
物种	Mouse
表达宿主	Human Cells
序列信息	Gln24-Asn148
检索号	P10148
分子量	14.7 kDa
标签	C-6His

### 产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	<1.0 EU per µg as determined by LAL test.
保存	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
运输	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
制剂	Supplied as a 0.2 µm filtered solution of PBS, pH 7.4.



## 背景介绍

C-C Motif Chemokine 2 (CCL2) is a member of the C-C or  $\beta$  Chemokine family. Mouse CCL2 shares 82% amino acid (aa) identity with rat CCL2 over the entire sequence, and 58%, 56%, 55%, 53% and 53% aa identity with human, equine, porcine, bovine and canine CCL2, respectively. Fibroblasts, glioma cells, smooth muscle cells, endothelial cells, lymphocytes and mononuclear phagocytes can produce CCL2 either constitutively or upon mitogenic stimulation, but monocytes and macrophages appear to be the major source. In addition to its chemotactic activity, CCL2 induces enzyme and Cytokine release by monocytes, NK cells and lymphocytes, and histamine release by basophils that express its Receptor, CCR2. Additionally, it promotes Th2 polarization in CD4+ T cells. CCL2-mediated recruitment of monocytes to sites of inflammation is proposed to play a role in the pathology of atherosclerosis, multiple sclerosis and allergic asthma.

## SDS-PAGE

